

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 5/00, A01N 63/00	A1	(11) International Publication Number: WO 96/23058 (43) International Publication Date: 1 August 1996 (01.08.96)
(21) International Application Number: PCT/US96/00900 (22) International Filing Date: 22 January 1996 (22.01.96) (30) Priority Data: 08/377,771 24 January 1995 (24.01.95) US (71) Applicant: CASE WESTERN RESERVE UNIVERSITY [US/US]; 10900 Euclid Avenue, Cleveland, OH 44106 (US). (72) Inventors: HAYNESWORTH, Stephen, E.; 3643 Antisdale Road, Cleveland Heights, OH 44118 (US). CAPLAN, Arnold, I.; 1300 Oak Ridge Drive, Cleveland Heights, OH 44121 (US). GERSON, Stanton, L.; 2463 Snowberry Lane, Pepper Pike, OH 44124 (US). LAZARUS, Hillard, M.; 2704 Rochester Road, Shaker Heights, OH 44122 (US). (74) Agents: HERRON, Charles, J. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).		(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ENHANCING BONE MARROW ENGRAFTMENT USING MSCs		
(57) Abstract Method and preparations for enhancing bone marrow engraftment in an individual by administering (i) a culturally expanded mesenchymal stem cell preparation and (ii) a bone marrow graft. The mesenchymal stem cells are administered in an amount effective to promote engraftment of the bone marrow.		

ENHANCING BONE MARROW ENGRAFTMENT USING MSCs

The present invention is directed to various methods and devices for using mesenchymal stem cells (MSCs) ~~to enhance~~ bone marrow engraftment. Mesenchymal stem cells are the formative pluripotent blast cells found in the bone that are capable of differentiating into any of the specific types of connective tissues (i.e., the tissues of the adipose, areolar, osseous, cartilaginous, elastic, and fibrous connective tissues) depending upon various environmental influences. Although these cells are normally present at very low frequencies in bone marrow and other mesenchymal tissues, the inventors of the present invention have discovered a process for isolating, purifying, and greatly replicating the mesenchymal stem cells in culture, i.e. in vitro. This discovery is the subject of co-pending U.S. patent application Serial No. 193,262, filed February 8, 1994. The methods and devices of the invention utilize such isolated and culture-expanded mesenchymal stem cells (MSCs). Under selected conditions, they can be induced to differentiate into different types of skeletal and connective tissues such as bone, cartilage, tendon, ligament, muscle, other connective tissues and marrow stroma.

The marrow stroma provides the scaffolding as well as soluble factors which direct and support blood cell

synthesis, i.e., hematopoiesis. The present invention is directed to a method to improve the process of bone marrow transplantation which is used to regenerate blood cells and marrow tissue in patients where their marrow is depleted or destroyed, such as, for example, during intensive radiation and chemotherapy treatment. Along this line, the inventors have discovered that, after lethal doses of radiation, culture-expanded MSCs can increase survival and decrease the time of blood and marrow cell regeneration when transplanted with complete bone marrow.

The present invention provides a method for enhancing the regeneration of marrow tissue through improved bone marrow transplantation using MSCs. The method for enhancing bone marrow engraftment comprises administering to an individual in need thereof, (i) mesenchymal stem cells and (ii) a bone marrow graft, wherein said mesenchymal stem cells are administered in an amount effective to promote engraftment of the bone marrow in the individual. More particularly, one embodiment of the invention is directed to a method for using a culture medium comprised of Dulbecco's Modified Essential Medium with low glucose (DMEM-LG) or medium 199 plus 1% human albumin as a vehicle or carrier for MSCs which, when administered systemically, will migrate, or home, to the marrow cavity and differentiate into marrow stroma, thereby regenerating the marrow stroma. The MSCs can be administered systemically, e.g., intravenously, into various delivery sites or directly into the bone.

A further aspect of the present invention is directed to the timing of injection of the MSCs into the patient relative to the bone marrow transplantation. In one embodiment, the MSCs are injected simultaneously with the bone marrow. In another embodiment, the MSCs are administered before or after the injection of bone marrow.

The present invention is useful to enhance the effectiveness of bone marrow transplantation as a treatment for cancer. The treatment of cancer by x-irradiation or alkylating therapy destroys the bone marrow microenvironment as well as the hematopoietic stem cells. The current treatment is to transplant the patient after marrow ablation with bone marrow which has been previously harvested and cryopreserved. However, because the bone marrow microenvironment is destroyed, bone marrow engraftment is delayed until the stromal environment is restored. As a result, a critical aspect of the present invention is directed to the advantages of transplanting isolated, purified, culture-expanded MSCs to accelerate the process of stromal reconstitution and ultimately marrow engraftment.

Modes of administration of the MSC preparation include but are not limited to systemic intravenous injection and injection directly to the intended site of activity. The preparation can be administered by any convenient route, for example by infusion or bolus injection and can be administered together with other biologically active agents. Administration is preferably systemic.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of the MSCs, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to medium 199 plus 1% serum albumin, saline, buffered saline, dextrose, water, and combinations thereof. The formulation should suit the mode of administration.

In a preferred embodiment, the MSC preparation or composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically,

compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a local anesthetic to ameliorate any pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a cryopreserved concentrate in a hermetically sealed container such as an ampoule indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The method of the invention can be altered, particularly by (1) increasing or decreasing the time interval between injecting MSCs and implanting the tissue; (2) increasing or decreasing the amount of MSCs injected; (3) varying the number of MSC injections; (4) varying the method of delivery of MSC; or (5) varying the source of MSCs. Although MSCs derived from the tissue donor is preferable, the MSCs can be obtained from other individuals or species, or from genetically-engineered inbred donor strains, or from in vitro cell culture.

The MSC preparation is used in an amount effective to promote engraftment of bone marrow in the recipient. In general, such amount is at least 1×10^4 MSC per kg of body weight and most generally need not be more than 7×10^5 MSC/kg. Preferably, it is at least about 2×10^5 MSC/kg prior to graft introduction and usually need not be more than about 7×10^5 MSC/kg. The MSC preparation may be administered concurrently with the bone marrow transplant or for a period prior to graft introduction of at least about 7 days but

generally not to exceed 30 days, with a typical therapeutic treatment period of 7 to 14 days. The MSC preparation preferably is administered either intravenously one to three times per day, and may be adjusted to meet optimal efficacy and pharmacological dosing.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical preparation of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

EXAMPLE 1

Isolation, Purification and Cultural

Expansion of Marrow-Derived Mesenchymal Stem Cells

Marrow in femoral head cancellous bone pieces was obtained from patients with degenerative joint disease during hip or knee joint replacement surgery. In addition, marrow was also obtained by iliac aspirate from normal donors and oncology patients who were having marrow harvested for future bone marrow transplantation. All of the oncology patients had malignancies unrelated to the stromal cells and the stromal cells expressed normal karyotype.

Marrow for cell culture was prepared from plugs of cancellous bone marrow as follows. Plugs of cancellous bone marrow (0.5-1.5 ml) were transferred to sterile tubes to which 25 ml Dulbecco's Modified Eagles Medium with low glucose (DMEM-LG) (GIBCO, Grand Island, N.Y.) with selected

batches of 10% fetal bovine serum (complete medium) was added. The tubes were vortexed to disperse the marrow, then spun at 1000 rpm for 5 minutes to pellet cells and bone pieces. The supernatant and fat layer were removed and the marrow and bone were reconstituted in 5 ml complete medium and vortexed to suspend the marrow cells. The suspended cells were collected with a 10 ml syringe fitted with a 16 gauge needle and transferred to separate tubes. Bone pieces were reconstituted in 5 ml of complete medium and the marrow cells were collected as before. Collection of marrow cells was considered complete when a pellet of yellowish-white cancellous bone pieces was all that remained in the original tube. Marrow cells were separated into a single cell suspension by passing them through syringes fitted with 18 and 20 gauge needles. Cells were spun at 1000 x g for 5 minutes after which the fat layer and supernatant were removed. Cells were reconstituted in complete medium, counted with a hemocytometer (red blood cells were lysed prior to counting with 4% acetic acid), and plated in 100 mm dishes at 50×10^6 nucleated cells/dish.

Marrow for cell culture was prepared from aspirated bone marrow as follows. Aspirate marrow (5-10 ml) was transferred to sterile 50 ml plastic centrifuge tubes to which 20 ml complete medium was added. The tubes were spun at 1,500 rpm for 5 minutes to pellet the cells. The supernatant and fat layer were removed and the cell pellets were resuspended to 5 ml with complete medium. The cell suspensions then were loaded onto 70% Percoll (Sigma, St. Louis, Mo.) gradients with a 10 ml pipette and spun at 2,000 rpm (460xg) in a GS34 rotor for 15 minutes. In order to harvest the cells, the tube is marked just below the high concentrated band of platelets, about 25% to 35% of the way down the tube. (Pooled density = 1.03g/ml.) Using a 10 ml pipette, the medium is aspirated off from the top down to the marked line

(approximately 12-14 ml). the collected fraction is transferred to a 50 ml conical plastic tube. 30 ml of complete medium then is added to the tube and centrifuged for 5 minutes at 1,500 rpm. The supernatant then is removed and discarded.

The cells then are resuspended in 7 ml of complete medium, and a uniform cell suspension is generated by pipetting cells up and down with a 10 ml pipette. The cells are counted by taking 20 μ l of uniform cell suspension, and adding 20 μ l of 4% acetic acid to the cell suspension. The suspension is mixed and transferred to a hemocytometer, and the cells are counted.

Cell concentration then is adjusted with complete medium to 50×10^6 cells per 7 ml. 7 ml of cells (50×10^6 cells) then are plated per 100mm culture plate.

The marrow mesenchymal stem cells were purified and expanded as follows. Marrow cells from either the femoral head cancellous bone or the iliac aspirate were cultured in complete medium (i.e., DMEM-LG medium with 10% fetal bovine serum) at 37°C in humidified atmosphere containing 95% air and 5% CO₂. On day 3 after plating, nonadherent cells were removed from the cultures by aspirating the original medium from the plates and replacing the original medium with 7 ml of fresh complete medium. Subsequent medium changes were performed every 3 to 4 days. This process of removing the non-adherent cells during culture media changes results in purification of the mesenchymal stem cells which selectively adhere to the culture plates. When primary culture dishes became nearly confluent, the medium was removed and each plate rinsed with 7 ml of sterile Tyrode's solution. The cells were detached with 4 ml of 0.25% trypsin with 0.1 mM EDTA (GIBCO) for 5 minutes at 37°C. The action of trypsin

was stopped with 2 ml of calf serum. The cells then were collected with a 10 ml pipette and centrifuged for 5 minutes at 1,500 rpm. The supernatant then is removed, and the cells resuspended from each plate in 21 ml complete medium, and a uniform cell suspension is generated by gently pipetting the cells up and down in the pipette. The cells from each culture then are counted and plated in three new 100 ml plates in 7 ml complete medium. The three new plates of cells are termed first passage cultures. First passage cells then were allowed to expand until they became near confluent, and were then replated as described above to create second passage cultures. At each passage, aliquots of cells were cryopreserved in 90% fetal bovine serum with 10% DMSO (freezing medium).

EXAMPLE 2

Mesenchymal stem cells were obtained from 10ml of autologous human bone marrow collected either at the time of autologous bone marrow harvest, or during a routine diagnostic bone marrow examination, from the iliac crest. These mesenchymal stem cells were culture expanded and infused into 14 patients who had hematologic malignancies, but had no evidence of active malignancy, and had not received chemotherapy or radiation therapy for at least 4 weeks before collection of mesenchymal stem cells. Five patients each received 1 million autologous mesenchymal stem cells, five patients each were given 10 million mesenchymal stem cells, and four patients each received 50 million mesenchymal stem cells. Toxicity for each patient was determined according to the National Cancer Institute toxicity grading scale. The toxicity grading scale was the same as that based upon the common toxicity criteria for a Phase II trial of high-dose sequential chemotherapy and peripheral blood stem cell autologous transplantation as initial therapy for patients with poor prognosis non-

Hodgkin's lymphoma. (National Cancer Institute document, draft 4/94: E2493.) Toxicity was graded on a scale from 0 to 4. No Grade 2 or greater toxic effects were observed, and in two patients that received 10 million mesenchymal stem cells, there appeared to be an increase in bone marrow cellularity.

EXAMPLE 3

Administration of Bone Marrow Cells and Mesenchymal Stem Cells to Breast Cancer Patients Treated with Chemotherapy

A breast cancer patient undergoes a diagnostic posterior iliac crest bone marrow aspiration and biopsy using a local anesthetic. A small portion (2 to 3 ml) of the aliquot (10 to 20 ml) of marrow is submitted for routine histologic testing and determination of tumor content using immunoperoxidase testing of cells grown in semisolid agar as described in Ross, et al., Blood, Vol. 82, pgs. 2605-2610 (1993). The remainder of the cells are cultured as hereinabove described. After at least 3 weeks in culture, samples of the mesenchymal stem cells are submitted for cell counting, cell viability (Trypan Blue), and cell surface immunophenotyping (testing with monoclonal antibodies SH2, SH3, and SH4). SH2 is produced by a hybridoma cell line assigned ATCC Accession No. HB10743. SH3 is produced by a hybridoma cell line assigned ATCC Accession No. HB10744. SH4 is produced by a hybridoma cell line assigned ATCC Accession No. HB10745. Samples also are submitted for detection of occult breast cancer.

The patient also undergoes placement of a pheresis central venous catheter, and receives subcutaneous injections of G-CSF (filgrastin) 10µg/kg/day as described in Peters, et al., Blood, Vol. 81, pgs. 1709-1719 (1993); Chao, et al.,

Blood, Vol. 81, pgs. 2031-2035 (1993); Sheridan, et al., The Lancet, Vol. 2, pgs. 891-895 (1989); and Winter, et al., Blood, Vol. 82, pg. 293a (1993). G-CSF injections begin at least 3 days before the first pheresis is initiated. G-CSF therapy is withheld if the white blood cell count rises above 40,000/ μ L and is resumed once the white blood cell count drops to less than 20,000/ μ L.

If the patient is receiving only G-CSF as the vehicle for "mobilization" of peripheral blood progenitor cells, the patient must not have received chemotherapy within 4 weeks of the planned pheresis. If the patient has received both conventional chemotherapy and G-CSF treatment for mobilization, the patient must not have received chemotherapy within 10 days of the planned pheresis, and the white blood cell count must be at least 800/ μ L and the platelet count at least 30,000/ μ L.

Daily pheresis procedures are performed using a Cobe Spectra instrument (Cobe, Lakewood, Colorado), and each cellular collection is cryopreserved using a controlled-rate liquid nitrogen freezer, until at least 15×10^8 mononuclear cells/kg are collected - (Lazarus, et al., Bone Marrow Transplant, Vol. 7, pgs. 241-246 (1991)). Each peripheral blood progenitor cell will be processed and cryopreserved according to previously published techniques. (Lazarus, et al., J. Clin. Oncol., Vol. 10, pgs. 1682-1689) (1992); Lazarus et al., (1991)).

Eight days before the patient is infused with the autologous peripheral blood progenitor cells, the patient receives chemotherapy over a period of 96 hours (4 days), with the following chemotherapy agents:

1. Cyclophosphamide in a total dosage of 6g/m^2 ($1.5\text{ g/m}^2/\text{day}$ for 4 days) is given via continuous intravenous infusion at 500 mg/m^2 in 1,000 ml normal saline every 8 hours.
2. Thiotepa in a total dosage of $500\text{ mg/m}^2/\text{day}$ for 4 days) is given via continuous intravenous infusion at 125 mg/m^2 in 1,000 ml normal saline every 24 hours.
3. Carboplatin in a total dosage of $1,800\text{ mg/m}^2$ ($200\text{ mg/m}^2/\text{day}$ for 4 days) is given via continuous intravenous infusion at 200 mg/m^2 in 1,000 ml of 5% dextrose in water every 24 hours.

The patient also receives 500 mg of mesna in 50 ml normal saline IV over 15 minutes every 4 hours for 6 days (144 hours), beginning with the first dose of cyclophosphamide.

At least 72 hours after the completion of the chemotherapy, the mesenchymal stem cells are harvested from tissue culture flasks. Cells are collected using bovine trypsin (which is deactivated with soybean trypsin inhibitor), suspended at 10×10^6 cells/ml in TC199 containing 1% human serum albumin (infusion medium), washed twice in infusion medium, resuspended at approximately 10^6 cells/ml, and injected slowly intravenously over 15 minutes to provide a total dosage of from 10 to about 5×10^6 cells.

The day after the patient receives the mesenchymal stem cells, the frozen autologous peripheral blood progenitor cells are removed from the liquid nitrogen refrigerator, transported to the patient in liquid nitrogen, submersed in a 37°C to 40°C sterile water bath, and infused rapidly intravenously without additional filtering or washing steps. GM-CSF in an amount of $250\text{ }\mu\text{g/m}^2$ then is given as a daily subcutaneous injection, beginning 3 hours after completion of the autologous blood progenitor cell infusion. The GM-CSF is

given daily until the peripheral blood neutrophil count exceeds 1,000/ μ L for three consecutive days.

It is to be understood, however, that the scope of the present invention is not to be limited to the specific embodiments described above. The invention may be practiced other than as particularly described and still be within the scope of the accompanying claims.

WHAT IS CLAIMED IS:

1. A method for enhancing bone marrow engraftment in an individual in need thereof which comprises administering to said individual (i) mesenchymal stem cells and (ii) a bone marrow graft, wherein said mesenchymal stem cells are administered in an amount effective to promote engraftment of said bone marrow in said individual.
2. The method of claim 1 wherein the mesenchymal stem cells are administered systemically.
3. The method of claim 1 wherein the preparation is administered intravenously.
4. The method of claim 1 wherein the preparation is injected into said cavity.
5. The method of claim 1 wherein the mesenchymal stem cells are autologous.
6. The method of claim 1 wherein the mesenchymal stem cell preparation is administered concurrently with the bone marrow graft.
7. The method of claim 6 wherein the mesenchymal stem cells are introduced into said individual in a cell suspension also containing bone marrow graft cells.
8. The method of claim 1 wherein the mesenchymal stem cell preparation is administered to the individual prior to administration of the bone marrow graft.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/00900

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 5/00; A01N 63/00

US CL : 435/240.2, 240.21; 424/93.1, 93.7

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/240.2, 240.21; 424/93.1, 93.7

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG

search terms: mesenchymal stem cells, bone marrow graft, bone marrow engraftment

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, E	US, A, 5486,359 (CAPLAN et al.) 23 January 1996, see entire document.	1-8
A	US, A, 5,226,914 (CAPLAN et al.) 13 July 1993, see entire document.	1-8
A	US, A, 5,197,985 (CAPLAN et al.) 30 March 1993, see entire document.	1-8



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
* A document defining the general state of the art which is not considered to be part of particular relevance	* X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E earlier document published on or after the international filing date	* Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* A document member of the same patent family
* O document referring to an oral disclosure, use, exhibition or other means	
* P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MARCH 1996

Date of mailing of the international search report

17 APR 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

Susan M. Dadio

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam